

## Access Full Complete Solution Manual Here

### Solution to problem 2.1

The first step is to calculate the equilibrium constant at 723 K. This constant can be determined from the standard free energy of reaction, as given by:

$$\Delta G_R^\circ = -R_g T \ln K$$

First calculate  $\Delta G_R^\circ$  at 298 K from  $\Delta G_f^\circ$  data (at 298 K). The following data are taken from the Appendix 1. The constants  $a$ ,  $b$ ,  $c$  and  $d$  are constants in the heat capacity polynomial.

Formula	$\Delta H_f^\circ$ kJ/mol	$\Delta G_f^\circ$ kJ/mol	$a$	$b \times 10^2$	$c \times 10^5$	$d \times 10^9$
N <sub>2</sub>	0	0	28.85	-0.1569	0.8067	-2.868
H <sub>2</sub>	0	0	29.06	-0.1913	0.3997	-0.8690
NH <sub>3</sub>	-46.22	-16.6	27.524	2.5603	0.98911	-6.6801

The free energy of formation of NH<sub>3</sub> is -16.6 kJ/mol. This value is equal to  $\Delta G_R^\circ$  because the free energies of formation of nitrogen and hydrogen are equal to zero. Therefore, at 298 K:

$$K_{298} = \exp\left(\frac{-\Delta G_R^\circ}{R_g T}\right) = \exp\left(\frac{-16.6 \times 10^3}{8.314 \times 298}\right) = 812$$


To calculate  $K$  at 723 K use the relationship:

$$\frac{\partial \ln K}{\partial T} = \frac{\Delta H_R^\circ}{R_g T^2}$$

This equation requires an expression for  $\Delta H_R^\circ$  as a function of  $T$ , which in turn depends on the difference in heat capacities between reactants and products, according to the equation:

$$\frac{\partial \Delta H_R^\circ}{\partial T} = \Delta C_P$$

The calculations are based on a reaction stoichiometry of one mole of NH<sub>3</sub>.



Using the data from Appendix 1 (given above), we can obtain an expression for the change in heat capacity. The general formula is:

$$\Delta C_P = \Delta a + \Delta bT + \Delta cT^2 + \Delta dT^3$$

Substituting the values from the above table gives:

$$\Delta C_P = -30.49 + 2.926 \times 10^{-2}T + 1.889 \times 10^{-5}T^2 - 3.943 \times 10^{-9}T^3$$

The enthalpy of reaction at 298 K is equal to the enthalpy of formation of  $\text{NH}_3$ , or  $\Delta H_{R,298}^\circ = -46.22 \times 10^3$  J/mol. By integration we then obtain the general equation:

$$\Delta H_R^\circ = \Delta H_{R,298}^\circ + \int_{298}^T \Delta C_P dT$$

Substitution and integration gives the following equation:

$$\Delta H_R^\circ = -46.22 \times 10^3 + \left[ -30.49T + \frac{2.926 \times 10^{-2}}{2}T^2 + \frac{1.889 \times 10^{-5}}{3}T^3 - \frac{3.943 \times 10^{-9}}{4}T^4 \right]_{298}^T$$

Simplify to give the following equation:

$$\Delta H_R^\circ = -3.86 \times 10^4 - 30.49T + 1.463 \times 10^{-3}T^2 + 6.297 \times 10^{-6}T^3 - 9.86 \times 10^{-10}T^4$$

The equilibrium constant at 723 K can now be computed:

$$\ln \left( \frac{K_{723}}{K_{298}} \right) = \frac{1}{R_g} \int_{298}^{723} \frac{\Delta H_R^\circ}{T^2} dT$$

Substitution and integration gives:

$$\ln \left( \frac{K_{723}}{812} \right) = \frac{1}{8.314} \left[ \frac{3.86 \times 10^4}{T} - 30.49 \ln(T) + 1.463 \times 10^{-3}T + \frac{6.297 \times 10^{-6}}{2}T^2 - \frac{9.86 \times 10^{-10}}{3}T^3 \right]_{298}^{723}$$

Substitution of the limits and simplification gives  $K_{723} = 4.153 \times 10^{-3}$ . The equilibrium constant can be expressed in terms of the mole fractions, fugacity coefficients and total pressure:

$$K = \frac{K_\phi K_y}{K_{f^\circ}} P^{\sum \nu_j}$$

where  $K_y = \frac{y_{\text{NH}_3}}{y_{\text{N}_2}^{0.5} y_{\text{H}_2}^{1.5}}$  and  $K_\phi = \frac{\phi_{\text{NH}_3}}{\phi_{\text{N}_2}^{0.5} \phi_{\text{H}_2}^{1.5}}$ .

The sum of the stoichiometric coefficients is:

$$\sum v_i = 1 - \frac{3}{2} - \frac{1}{2} = -1$$

The fugacity coefficient  $\phi$  is defined as:

$$\phi = \frac{f}{P}$$

Therefore the three fugacity coefficients are:

$$\phi_{\text{H}_2} = \frac{1350}{1000} = 1.35, \quad \phi_{\text{N}_2} = \frac{1380}{1000} = 1.38 \quad \text{and} \quad \phi_{\text{NH}_3} = \frac{860}{1000} = 0.86$$

Therefore:

$$K_\phi = \frac{0.86}{(1.38)^{0.5} (1.35)^{1.5}} = 0.467$$

The mole fractions in  $K_y$  refer to equilibrium mole fractions. Take a basis of 100 moles of feed and let  $w$  moles of  $\text{N}_2$  react. The following mole table then can be constructed:

Component	Initial Moles	Equilibrium Moles
$\text{N}_2$	21	$21 - w$
$\text{H}_2$	63	$63 - 3w = 3(21-w)$
$\text{NH}_3$	0	$2w$
Ar	16	16
Total	100	$100 - 2w$

Express  $K_y$  in terms of equilibrium moles.

$$K_y = \left( \frac{2w}{100 - 2w} \right) \left( \frac{100 - 2w}{3(21 - w)} \right)^{1.5} \left( \frac{100 - 2w}{21 - w} \right)^{0.5}$$

Now substitution of  $K$ ,  $K_\phi$  and  $P$  gives:

$$K = K_y K_\phi P^{-1} \quad \text{or} \quad 4.153 \times 10^{-3} = K_y (0.467) (1000)^{-1}$$

Solving gives  $K_y = 8.893$ . The value of  $w$  can then be determined. Simplification gives a quadratic equation:

$$w^2 - 42.64w + 405.9 = 0$$

This equation can be solved using the quadratic formula. It has two roots,  $w = 15.36$  and  $w = 27.28$ . As  $w$  must be less than 21, it follows that the realistic solution is  $w = 15.36$ . Therefore, the equilibrium mole fractions at 1000 atm and 723 K are:

$$N_2 = \frac{21 - 15.36}{100 - 2 \times 15.36} = 0.0814$$

$$H_2 = \frac{3(21 - 15.36)}{100 - 2 \times 15.36} = 0.244$$

$$NH_3 = \frac{2 \times 15.36}{100 - 2 \times 15.36} = 0.443$$

$$Ar = \frac{16}{100 - 2 \times 15.36} = 0.231$$

**Comments:** High pressure favours conversion because there is a decrease in moles on reaction. The reaction is exothermic and therefore a low temperature would give a more favourable conversion. However, a high temperature is used to give a high reaction rate.

### Solution to problem 2.2

#### Part (a)

The first step is to calculate the equilibrium constant at 1023 K. For this purpose the equilibrium constant is first calculated at 298 K using the free energy of reaction:

$$\Delta G_R^\circ = -R_g T \ln K$$

$K$  is then calculated at 1023 K using the van't Hoff equation:

$$\frac{\partial \ln K}{\partial T} = \frac{\Delta H_R^\circ}{R_g T^2}$$

The Gibbs free energy of reaction is calculated from the data in Appendix 1. The data, including heat capacity data for the species present are:

Formula	$\Delta H_f^\circ$ kJ/mol	$\Delta G_f^\circ$ kJ/mol	$a$	$b \times 10^2$	$c \times 10^5$	$d \times 10^9$
H <sub>2</sub>	0	0	29.06	-0.1913	0.3997	-0.8690
C <sub>2</sub> H <sub>6</sub>	-84.72	-32.9	6.889	17.24	-6.395	7.273
C <sub>2</sub> H <sub>4</sub>	52.32	68.17	3.95	15.61	-8.331	17.64

Therefore the free energy of reaction at 298 K is:

$$\begin{aligned} \Delta G_{R,298}^\circ &= \Delta G_{f,298}^\circ(\text{C}_2\text{H}_4) - \Delta G_{f,298}^\circ(\text{C}_2\text{H}_6) \\ \Delta G_{R,298}^\circ &= 68.05 \times 10^3 - (-32.90 \times 10^3) = 100.95 \times 10^3 \frac{\text{J}}{\text{mol}} \end{aligned}$$

The equilibrium constant at 298 K is then calculated to be  $K_{298} = 1.9373 \times 10^{-18}$ . To compute  $K_{1023}$  it is first necessary to calculate  $\Delta H_R^\circ$  as a function of temperature. For this purpose use the following relationship:

$$\frac{\partial \Delta H_R^\circ}{\partial T} = \Delta C_P$$

The difference in heat capacity between products and reactants is calculated from the above data. The general formula is:

$$\Delta C_P = \Delta a + \Delta bT + \Delta cT^2 + \Delta dT^3$$

Substituting the values from the above table gives:

$$\Delta C_P = 25.5 - 1.222 \times 10^{-2} T - 2.788 \times 10^{-5} T^2 + 16.45 \times 10^{-9} T^3$$

The enthalpy change with reaction at 298 K is:

$$\Delta H_{R,298}^\circ = \Delta H_{f,298}^\circ(\text{C}_2\text{H}_4) - \Delta H_{f,298}^\circ(\text{C}_2\text{H}_6)$$

$$\Delta H_{R,298}^\circ = 52.25 \times 10^3 - (-84.60 \times 10^3) = 136.9 \times 10^3 \frac{\text{J}}{\text{mol}}$$

The general expression for  $\Delta H_R^\circ$  is thus derived by integration:

$$\Delta H_R^\circ = \Delta H_{R,298}^\circ + \int_{298}^T \Delta C_P dT$$

For the cracking of ethane, substitution of the values and integration gives:

$$\Delta H_R^\circ = 130,007 + 25.5 T - 6.11 \times 10^3 T^2 - 9.293 \times 10^{-6} T^3 + 4.113 \times 10^{-9} T^4$$

The general equation for the equilibrium constant in integral form is:

$$\ln K = \ln K_{298} + \frac{1}{R_g} \int_{298}^T \frac{\Delta H_R^\circ}{T} dT$$

Substitution of  $\Delta H_R^\circ$  and integration with a limit of 1023 K gives a value of  $K_{1023} = 0.4916$ .

The equilibrium constant is related to the composition by:

$$K = \frac{K_\phi K_y}{K_{f^\circ}} P^{\sum v_j}$$

where 
$$K_y = \frac{y_{\text{H}_2} y_{\text{C}_2\text{H}_4}}{y_{\text{C}_2\text{H}_6}} \quad \text{and} \quad K_\phi = \frac{\phi_{\text{H}_2} \phi_{\text{C}_2\text{H}_4}}{\phi_{\text{C}_2\text{H}_6}} = 1.$$

The sum of the stoichiometric coefficients is:

$$\sum v_i = 1 + 1 - 1 = 1$$

Therefore:

$$K = K_y P = \frac{y_{\text{H}_2} y_{\text{C}_2\text{H}_4}}{y_{\text{C}_2\text{H}_6}} P$$

with  $P$  in atm.  $P = 120/100 = 1.2$  bar. Take a basis of 1 mole of ethane and let  $X$  moles react ( $X$  is therefore the fractional conversion).

Compound	Initial Moles	Final Moles
$C_2H_6$	1	$1 - X$
$C_2H_4$	0	$X$
$H_2$	0	$X$
Total	1	$1 + X$

Substitute for the mole fractions:

$$K_y = \left( \frac{X}{1+X} \right) \left( \frac{X}{1+X} \right) \left( \frac{1+X}{1-X} \right) = \frac{X^2}{(1-X)(1+X)}$$

Therefore, the equilibrium conversion is given by:

$$0.4916 = \frac{X^2}{(1-X^2)} 1.2$$

Solve for  $X = 0.539$ . The conversion of ethane is 53.9 % at equilibrium.

### Part (b)

In part (a) a general expression for  $\Delta H_R^\circ$  was derived. Substituting 1023 K into this equation gives  $\Delta H_R^\circ = 144255$  J/mol. This number is positive which indicates that 144,255 J must be transferred to the reactor for each mole of ethane that cracks.

### Solution to problem 2.3

The equilibrium constant at 500 °C has a value of 85. The relationship between composition and the equilibrium constant  $K$  is:

$$K = \frac{K_y K_\phi}{K_{f_i^\circ}} P^{\sum v_i}$$

From the reaction stoichiometry the change in moles on reaction is:

$$\sum v_i = 1 - 1 - \frac{1}{2} = -\frac{1}{2}$$

Because the fugacity coefficients are equal to one, it follows that  $K_\phi = 1$ . Using a standard state of 1 bar gives:

$$K = K_y P^{-0.5} \quad \text{where } P \text{ is in bar}$$

The mole fractions at equilibrium must be expressed in terms of the number of moles present at equilibrium. Take a basis of 100 moles of original mixture and let  $X$  be the fractional conversion at equilibrium. That is,

$$X = \frac{N_{\text{SO}_2,0} - N_{\text{SO}_2}}{N_{\text{SO}_2,0}}$$

Construct a stoichiometric table on the 100 mole basis:

Compound	Moles Initial	Moles Final
SO <sub>2</sub>	8	8(1 - X)
SO <sub>3</sub>	0	8X
O <sub>2</sub>	11	11 - $\frac{8}{2}X$
N <sub>2</sub>	81	81
Total	100	100 - 4X